

# Results of the global conservation assessment of the freshwater crabs (Brachyura, Pseudothelphusidae and Trichodactylidae): The Neotropical region, with an update on diversity

Neil Cumberlidge<sup>1</sup>, Fernando Alvarez<sup>2</sup>, Jose-Luis Villalobos<sup>2</sup>

**1** *Department of Biology, Northern Michigan University, Marquette, MI 49855, USA* **2** *Colección Nacional de Crustáceos, Instituto de Biología, Universidad Nacional Autónoma de México, Apartado Postal 70–153, México 04510, D. F. México*

Corresponding author: Neil Cumberlidge ([ncumberl@nmu.edu](mailto:ncumberl@nmu.edu))

---

Academic editor: I. S. Wehrtmann | Received 12 November 2013 | Accepted 21 March 2014 | Published 25 November 2014

---

<http://zoobank.org/79468B34-97FF-41F6-B849-4B1B80D9D0AD>

---

**Citation:** Cumberlidge N, Alvarez F, Villalobos J-L (2014) Results of the global conservation assessment of the freshwater crabs (Brachyura, Pseudothelphusidae and Trichodactylidae): The Neotropical region, with an update on diversity. In: Wehrtmann IS, Bauer RT (Eds) Proceedings of the Summer Meeting of the Crustacean Society and the Latin American Association of Carcinology, Costa Rica, July 2013. ZooKeys 457: 133–157. doi: 10.3897/zookeys.457.6598

---

## Abstract

The freshwater crabs of the Neotropics comprise 311 species in two families (Pseudothelphusidae and Trichodactylidae) and one or both of these families are found in all of the countries in the Neotropical region (except for Chile and some of the Caribbean islands). Colombia (102 species, 81% endemic) and Mexico (67 species, 95% endemic) are the biodiversity hotspots of freshwater crab species richness and country-level endemism for this region. The results of the IUCN Red List conservation assessments show that 34% of pseudothelphusids and 10% of trichodactylids have an elevated risk of extinction, 29% of pseudothelphusids and 75% of trichodactylids are not at-risk (Least Concern), and although none are actually extinct, 56% of pseudothelphusids and 17% of trichodactylids are too poorly known to assess (Data Deficient). Colombia (14 species), Venezuela (7 species), Mexico (6 species), and Ecuador (5 species) are the countries with the highest number of threatened species of Neotropical freshwater crabs. The majority of threatened species are restricted-range semiterrestrial endemics living in habitats subjected to deforestation, alteration of drainage patterns, and pollution. This underlines the need to prioritize and develop conservation measures before species decline to levels from which they cannot recover. These results represent a baseline that can be used to design strategies to save threatened Neotropical species of freshwater crabs.

**Keywords**

Pseudothelphusidae, Trichodactylidae, Neotropical region, conservation, distribution, endemism

**Introduction**

The Neotropical region occupies the entire South American continent, plus Mexico and Central America, the islands of the Caribbean, and southern Florida. The Nearctic/Neotropical boundary runs through Mexico and passes through the southern parts of Baja California and Sonora, crosses the Meseta Central, and continues to southern Veracruz. The majority of the Neotropical region has a tropical climate with warm water freshwater ecosystems, but the southern part of this region from 10° to 25°S (southern Brazil, Uruguay, Paraguay, Argentina, and Chile) has a subtropical climate with cooler freshwater habitats. Freshwater crabs are found throughout the freshwater ecosystems of the Neotropical region but are notably absent from Baja California and the Yucatan Peninsula (Mexico), southern Florida (USA), some of the Caribbean islands, southern Argentina, and all of Chile.

Our knowledge of the impressive freshwater crab fauna in this region has been very slow to develop since the first species of trichodactylid and pseudothelphusid freshwater crabs were described in 1783 and 1840, respectively. Only one species (a trichodactylid) was described in the 18<sup>th</sup> C (Herbst 1783), 46 more species were described in the 19<sup>th</sup> C (Ortmann 1893, 1897; Rathbun 1893, 1898), 209 more species were added in the 20<sup>th</sup> C (Rathbun 1905, Smalley 1964a,b, 1965, 1970, Bott 1967, 1968, 1969, 1970, Rodríguez and Smalley 1969, Pretzmann 1972, 1975, Hobbs 1980, Rodríguez 1980, 1982, Campos and Rodríguez 1984, 1985, 1993, 1995, Smalley and Adkison 1984, Magalhães 1986, Magalhães and Türkay 1986, Rodríguez 1986, Alvarez 1987, 1989, Campos and Rodríguez 1988, Campos 1989, Alvarez and Villalobos 1990, 1991, 1994, 1995, 1996, 1997a,b, 1998, Hobbs 1991, Campos 1992, 1999, Rodríguez 1992, Villalobos et al. 1993, Campos 1995, 1997, Alvarez et al. 1996, 1999, Magalhães and Türkay 1996a–c, Campos and Lemaitre 1998), and 31 species have been described so far in the 21<sup>st</sup> C (Campos 2000, 2001, 2002, 2003a, b, 2004, 2005, 2010a, b, 2011, Campos and Lemaitre 2002, Rodríguez et al. 2002, Villalobos and Alvarez 2003, 2009, 2010, 2013, Campos and Valencia 2004, Rodríguez and Magalhães 2005, Villalobos 2005, Campos and Pedraza 2006, 2008, Cumberlidge 2007, Magalhães and Türkay 2008a, b, c, 2012, Ng et al. 2008, Yeo et al. 2008, Cumberlidge et al. 2009, Magalhães 2010, Magalhães and Türkay 2010, Magalhães et al. 2010, Villalobos et al. 2010, Alvarez et al. 2012a, Cardona and Campos 2012, Magalhães et al. 2013). By 2009, the total number of Neotropical species stood at more than 300, making this the second most species-rich region in the world after the Oriental region (Ng et al. 2008, Cumberlidge et al. 2009). Interest in the Neotropical freshwater crab fauna remains strong and 13 new species have already been recognized since 2009 (Table 1). Exploration is still continuing and there is every prospect that the species count will increase further as taxonomic discrimination improves. We expect most of

**Table 1.** Species of Neotropical freshwater crabs described since Cumberlidge et al. (2009).

Family	Subfamily/Tribe	Species	Country
Pseudothelphusidae	Kingsleyini	<i>Microthelphusa lipkei</i> Magalhães, 2010	Brazil
Pseudothelphusidae	Kingsleyini	<i>Brasiliothelphusa dardanelosensis</i> Magalhães & Türkay, 2010	Brazil
Pseudothelphusidae	Strengerianini	<i>Phallangothelphusa juansei</i> Campos, 2010	Colombia
Pseudothelphusidae	Strengerianini	<i>Phallangothelphusa martensis</i> Cardona & Campos, 2012	Colombia
Pseudothelphusidae	Hypolobocerini	<i>Neostrengeria alexae</i> Campos, 2010	Colombia
Pseudothelphusidae	Hypolobocerini	<i>Neostrengeria natashae</i> Campos, 2011	Colombia
Pseudothelphusidae	Potamocarcinini	<i>Potamocarcinus darienensis</i> Magalhães, Campos & Türkay, 2013	Panama
Pseudothelphusidae	Hypolobocerini	<i>Allacanthos yawi</i> Magalhães, Lara & Wehrtmann, 2010	Costa Rica
Pseudothelphusidae	Potamocarcinini	<i>Odontothelphusa apicpac</i> Villalobos, García & Velázquez, 2010	Mexico
Pseudothelphusidae	Potamocarcinini	<i>Sylvathelphusa kalebi</i> Villalobos & Alvarez, 2013	Mexico
Pseudothelphusidae	Potamocarcinini	<i>Sylvathelphusa cavernicola</i> Villalobos & Alvarez, 2013	Mexico
Pseudothelphusidae	Potamocarcinini	<i>Tzotzithelphusa villarosalensis</i> Villalobos & Alvarez, 2013	Mexico
Pseudothelphusidae	Pseudothelphusini	<i>Pseudothelphusa zongolicae</i> Alvarez, Villalobos & Moreno, 2012	Mexico

the new species to belong to the Pseudothelphusidae, because discovery of new species of trichodactylids has slowed dramatically in the last two decades.

Research on the Neotropical freshwater crabs has also focused on their phylogeny (Rodríguez and Campos 1989, 1998, Rodríguez and Pereira 1992, Rodríguez and Sternberg 1998, Sternberg et al. 1999, Sternberg and Cumberlidge 2001, Rodríguez and Magalhães 2005, Villalobos and Alvarez 2010), their biogeography (Rodríguez 1986, Campos 2005, Huidobro et al. 2006, Alvarez et al. 2012b, Ojeda et al. 2013), and their medical importance with 22 species of pseudothelphusids serving as second intermediate hosts of the human lung fluke, *Paragonimus* (Acha and Szyfres 2001) in 9 countries from Mexico and Central America, to the Andes, and Brazil (Rodríguez and Magalhães 2005).

The objectives of the present study are to describe and update our knowledge of Neotropical freshwater crab diversity of both the Pseudothelphusidae and Trichodactylidae and to describe the patterns of distribution and endemism within these families. We also identify here not only those species that are most vulnerable to extinction, but also those species that are poorly known and obvious candidates for future research attention.

## Methods

Comprehensive distributional data for the Neotropical freshwater crabs were compiled from literature and museum records, particularly the major monographs on the Pseudothelphusidae by Rodríguez (1982, 1992) and Rodríguez and Magalhães (2005), and

the Trichodactylidae by Rodríguez (1992) and Magalhães and Türkay (1996a–c, 2008a–b, 2012). Other regional monographs referred to include those for Colombia (Campos 2005), Ecuador (Rodríguez and Sternberg 1998), Brazil (Magalhães 2003), Peru (Rodríguez and Suárez 2004), and Mexico and Central America (Alvarez and Villalobos 1990, 1991, 1994, 1995, 1996, 1997, Villalobos et al. 1993, Alvarez et al. 1996, 1999, Villalobos and Alvarez 2003, 2008, 2009, 2010, Villalobos 2005). Despite these efforts distributional records for most species of freshwater crabs are still likely to be incomplete. Distribution maps presented here are based on specimen-level databases compiled for all species based on all known specimens and includes information from over 2,500 different localities. Individual conservation assessments for all species of Neotropical freshwater crabs assessed by Cumberlidge et al. (2009) using the IUCN Red List categories and criteria at the global scale (IUCN 2003) are provided on the IUCN Red List site (<http://www.iucn.redlist.org>). Freshwater crab species were assessed for inclusion in one of the Red List categories based on a combination of data on geographic range and/or population levels and related trends (Cumberlidge et al. 2009). The available data were sufficient to make valid assessments of the conservation status of only 150 out of the 298 species known from the Neotropical region at the time, and assessments were not possible for 148 species that were treated as Data Deficient due to a lack of specimens, and of locality and population data (Cumberlidge et al. 2009). Threats were inferred if a species was potentially subject to anthropogenic impacts such as habitat destruction, alteration of drainage patterns, or pollution, especially if it was either not found in a protected area, or if it was found in a protected area for only part of its range. Distribution maps (Figs 1–4) were prepared using ArcView GIS mapping software.

## Results

### Neotropical freshwater crab families

The Neotropical region has its own distinctly recognizable freshwater crab fauna, with no species occurring in other parts of the world. The Neotropical freshwater crab fauna is not uniformly distributed and species and genus composition changes from country to country (Table 2, Figs 1, 2). The global distributions of all species of the Pseudothelphusidae and Trichodactylidae based on all known localities are shown in figures 1 and 2. The Neotropical freshwater crab fauna today (49 genera, 311 species, 2 families) easily exceeds that of the entire Afrotropical region (18 genera, 145 species, 2 families), but is significantly less diverse than the Palaearctic-Oriental-Australasian region (147 genera, 850 species, 2 families) (Cumberlidge et al. 2009; present work).


The Neotropical freshwater crab fauna, which represent 20–22% of the world's freshwater crab species diversity (Ng et al. 2008, Yeo et al. 2008, Cumberlidge et al. 2009), is dominated by the highly diverse pseudothelphusids (34 genera, 264 species). Pseudothelphusids are found in 26 out of the 31 countries and reach their highest species richness in South America (Table 2). Pseudothelphusids are found throughout

**Table 2.** Distribution of Pseudothelphusidae (P) and Trichodactylidae (T) in each country in the Neotropical region that has populations of freshwater crabs. First number indicates the number of genera, species, and localities of Pseudothelphusidae, the second number indicates the same data for the Trichodactylidae.

	No. GENERA (P, T)	No. SPECIES (P, T)
<b>South America</b>		
Argentina	0, 5	0, 9
Bolivia	0, 6	0, 8
Brazil	6, 10	19, 31
Colombia	13, 8	88, 14
Ecuador	3, 4	21, 5
French Guiana	2, 4	3, 4
Guyana	3, 3	8, 3
Paraguay	0, 6	0, 7
Peru	2, 9	8, 14
Suriname	3, 3	5, 3
Uruguay	0, 1	0, 2
Venezuela	11, 7	35, 9
<b>Mexico and Central America</b>		
Mexico	15, 2	63, 4
Belize	4, 0	4, 0
Costa Rica	4, 0	14, 0
El Salvador	4, 0	6, 0
Guatemala	6, 0	12, 0
Honduras	2, 0	3, 0
Nicaragua	3, 3	5, 3
Panama	4, 1	14, 1
<b>Caribbean</b>		
Cuba	2, 0	9, 0
Dominica	1, 0	1, 0
Dominican Republic	1, 0	2, 0
Guadeloupe	1, 0	1, 0
Haiti	1, 0	1, 0
Martinique	1, 0	1, 0
Puerto Rico	1, 0	1, 0
Saint Vincent & the Grenadines	1, 0	1, 0
St Croix US Virgin Islands	1, 0	1, 0
Trinidad & Tobago	2, 1	2, 1

the warmer parts of the Neotropics from Mexico to Central America and the Andes (Venezuela, Colombia, Ecuador, and Peru), and in the Guyanian Shield and Central Brazilian Shield in the Amazon basin (Fig. 1, Table 2). The majority of pseudothelphusid species live at altitudes greater than 400 m above sea level, with a few Andean species reaching up to 3000 m (Campos 2005).


This family comprises two subfamilies, Epilobocerinae and Pseudothelphusinae (Ng et al. 2008). The Epilobocerinae consists of only 9 species in 2 genera (*Epilobocera*,




**Figure 1.** Distribution of Pseudothelphusidae based on all known point localities (n = 1719).

*Neopilobocera*) that are all endemic to the larger islands of the Greater Antilles (Cuba, Haiti, Dominican Republic, Puerto Rico, and St. Croix). The vast majority of pseudothelphusids are in the subfamily Pseudothelphusinae assigned to five Tribes: Pseudothelphusini (Mexico), Potamocarcinini (Central America and Mexico), Hypolobocerini (from Venezuela to Peru, plus Central America, and Mexico), Strengerianini (Colombia and Venezuela), and Kingsleyini (the Lesser Antilles, Venezuela, Colombia, and northern Brazil). The eastern limits of the Pseudothelphusidae in the Amazon basin is in Ceará State, Brazil, while their southwestern limit in the Amazon basin is marked by the watershed of the Ucayali River, south of which there are no species of pseudothelphusids (Rodríguez and Magalhães 2005).


The Trichodactylidae is the smallest of all freshwater crab families (15 genera, 47 species) and represents only 4% of the world's freshwater crab diversity. Despite this, the family is widely distributed and has representatives in 16 countries, most of which are in South America (Table 2) (Rodríguez 1992, Ng et al. 2008, Yeo et al. 2008, Cumberlidge et al. 2009). The vast majority of trichodactylid species are found below 100 m above sea level in the Magdalena river basin and Lake Maracaibo in Colombia and Venezuela, in the coastal lowlands of the Guyanas and Brazil, and in the lowland river basins of the Amazon, Orinoco, Paraguay, and Parana. A few species of trichodactylids reach up to 900 m above sea level in the Andes foothills (Campos 2005). Countries with trichodactylids include Colombia, Venezuela, French Guiana, Guyana, Suriname, Brazil, Ecuador, Peru, Bolivia, Paraguay, Uruguay, and Argentina (Fig. 2, Table 2). Outside of South America there are four trichodactylid species in Veracruz, Tabasco, Oaxaca, and Chiapas in Mexico



**Figure 2.** Distribution of Trichodactylidae based on all known point localities (n = 853).



**Figure 3.** Distribution of threatened species of Neotropical freshwater crabs (Pseudothelphusidae and Trichodactylidae) based on all known point localities (n = 173).



**Figure 4.** Distribution of Data Deficient species of Neotropical freshwater crabs (Pseudothelphusidae and Trichodactylidae) based on all known point localities (n = 338).

(*Avotrichodactylus constrictus*, *A. oaxensis*, *Rodriguezia mensabak*, and *R. villalobosi*) (Magalhães and Türkay 2012); one species (*Trichodactylus quinquedentatus*) from Colombia and Nicaragua (Smalley and Rodríguez 1972), and one species (*Poppiana dentata*) from Trinidad (which is also found on the neighboring mainland in Venezuela, Suriname, Guyana, and French Guiana, and in Nicaragua) (Magalhães and Türkay 2008a). The Trichodactylidae includes two subfamilies: the Trichodactylinae, with 3 genera, two from Mexico (*Avotrichodactylus* and *Rodriguezia*) and one from South and Central America (*Trichodactylus*); and the Dilocarcininae, with 9 genera that are found throughout South America (Rodríguez 1992, Magalhães and Türkay 1996a, b, c, 2008a, b). Interestingly, this classification (Ng et al. 2008) groups together at the subfamily level the four disjunct species of trichodactylids found in Mexico with the genus *Trichodactylus* that includes species from the Magdalena basin and Amazon slopes in Colombia, a species from Nicaragua and Colombia, species from the Atlantic coastal river systems of southeastern Brazil, and species from the Paraguay and Parana basins in southern Brazil and Argentina.

### Hotspots of freshwater crab diversity

Colombia stands out as the most diverse and the most species rich part of the region (21 genera, 102 species, 2 families) (Table 2) with biodiversity hotspots located in forested mountainous regions where deep valleys and a complex topographical relief are favorable for genetic isolation and allopatric speciation. The pseudothelphusid fauna of Colombia

comprises 13 genera and 88 species in one subfamily (Pseudothelphusinae) and four tribes (Hypolobocerini, Strengerianini, Kingsleyini, and Potamocarcinini). There are distinct northern and southern components of pseudothelphusid distribution in Colombia, with the dividing line coinciding with the watersheds of the San Juan, Cauca, and Magdalena river basins. The southern component of this fauna is dominated by *Hypolobocera*, *Lindacatalina*, and *Moritschus* and extends from southern Colombia into Ecuador and Peru, and also includes species from southeast Colombia and the Amazonian and Pacific slopes of Ecuador. The trichodactylid fauna of Colombia comprises 8 genera, 14 species in two subfamilies: Trichodactylinae (one genus) and Dilocarcininae (7 genera).

The next most species-rich country is Mexico (17 genera, 67 species, 2 families) (Table 2) with a freshwater crab fauna dominated by pseudothelphusids with 15 genera, and 63 species in one subfamily (Pseudothelphusinae) and three tribes (Pseudothelphusini, Potamocarcinini, and Hypolobocerini). Mexico's trichodactylid fauna is much smaller and comprises just 4 species in 2 genera all in the subfamily Trichodactylinae. Freshwater crab diversity in Mexico is highest in the Isthmus of Tehuantepec and Chiapas State, possibly because this is where species from three pseudothelphusid tribes overlap with species of trichodactylids (Villalobos and Alvarez 2010).

Brazil is the third most species-rich Neotropical country (16 genera, 50 species, 2 families) with a fauna that is dominated by trichodactylids (10 genera, 31 species, 2 subfamilies), and fewer pseudothelphusids (6 genera, 19 species, all Pseudothelphusinae, Kingsleyini). Venezuela is the fourth most species-rich Neotropical country (18 genera, 44 species, 2 families), made up of 11 genera and 35 species of pseudothelphusids (all Pseudothelphusinae, in three tribes, Hypolobocerini, Kingsleyini, and Strengerianini), and 7 genera and 9 species of trichodactylids (all Dilocarcininae).

The freshwater crab fauna of Ecuador (7 genera, 26 species, 2 families) includes 3 genera and 21 species of pseudothelphusids (all Pseudothelphusinae, Hypolobocerini), and 4 genera and 5 species of trichodactylids (all Dilocarcininae). Finally, Peru (11 genera, 22 species, 2 families) has a fauna dominated by trichodactylids (14 species and 8 genera, in 2 subfamilies Dilocarcininae and Trichodactylinae), with a smaller pseudothelphusid fauna consisting of 8 species and 2 genera (all Pseudothelphusinae in 2 tribes Kingsleyini and Hypolobocerini). Freshwater crabs of both families are completely absent from Chile. Despite warm subtropical climates freshwater crabs are absent in many islands of the Caribbean including Jamaica, The Cayman Islands, The Bahamas, Grenada, Antigua, and Anguilla.

When distribution patterns are considered at the genus level (Table 2) the taxonomic diversity of Neotropical freshwater crabs is again by far the highest in Colombia (21 genera), Venezuela (18 genera), Mexico (17 genera), Brazil (16 genera), and Peru (11 genera). Diversity is lower in Ecuador (7 genera), Guyana, Suriname, Bolivia, Paraguay, French Guiana, Nicaragua, and Guatemala (6 genera), and in Argentina and Panama (5 genera) (Table 2). Costa Rica, El Salvador and Belize each have 4 genera, Trinidad and Tobago has 3 genera, Honduras and Cuba each have 2 genera, while Uruguay and the other islands of the Greater and Lesser Antilles each have 1 genus.

### Widespread species: Pseudothelphusidae

Thirteen (out of 34) pseudothelphusid freshwater crab genera and 28 (out of 283) species have a wide distribution that easily exceeds 20,000 km<sup>2</sup> and includes more than one country. Notably, *Fredius reflexifrons* is found in six countries (Brazil, French Guiana, Guyana, Peru, Suriname, Venezuela), *Potamocarcinus magnus* is found in five countries (Mexico, Guatemala, Honduras, El Salvador, Costa Rica), and *Kingsleya latifrons* (Guyana, French Guiana, Suriname, Brazil) and *Raddaus bocourti* (Mexico, Belize, Guatemala, El Salvador) are found in four countries. Other widespread species found in three countries include *Potamocarcinus armatus* (Mexico, Guatemala, Nicaragua), *P. richmondi* (Nicaragua, Costa Rica, Panama), *Fredius denticulatus* (Brazil, French Guiana, Suriname) and *F. fittkaui* (Venezuela, Guyana, Brazil), and *Prionothelphusa eliasi* (Brazil, Colombia and Venezuela). Finally, 17 widespread species are found in two countries: *Rodriguezus garmani* (Trinidad and Tobago, Venezuela), *Fredius beccarii* (Guyana, Venezuela), *Zilchia zilchi* (El Salvador, Honduras), *Hypolobocera caputii* and *Moritschus henrici* (both Ecuador, Peru), *Ptychophallus exilipes* (Costa Rica, Panama), *Hypolobocera bouvieri angulata*, *Orthothelphusa holthuisi* (Colombia, Venezuela), and *Hypolobocera exuca*, *Lindacatalina latipenis*, *L. orientalis*, and *L. sumacensis* (all Colombia, Ecuador), *Fredius estevisi* and *F. platyacanthus* (Brazil, Venezuela), *Kingsleya siolii* (Brazil, Suriname), *Zilchia aspoekorum* (Belize, Mexico), and *Raddaus tuberculatus* (Guatemala, Mexico).

The majority of pseudothelphusids (22 species) that have a widespread distribution exceeding 20,000 km<sup>2</sup> are assessed as Least Concern. However, there are some species with a small distributional range that spans the border between two neighboring countries that are assessed as either Vulnerable: *Hypolobocera exuca* (Colombia, Ecuador), *Kingsleya siolii* (Brazil, Suriname) and *Zilchia aspoekorum* (Belize, Mexico), or Data Deficient: *Fredius platyacanthus* (Brazil, Venezuela), *Lindacatalina latipenis* and *L. sumacensis* (both Colombia, Ecuador), *Orthothelphusa holthuisi* (Colombia, Venezuela), and *Ptychophallus exilipes* (Costa Rica, Panama).

### Widespread Species: Trichodactylidae

Some 8 out of 15 trichodactylid freshwater crab genera and 20 out of 47 species have a wide distributional range exceeding 20,000 km<sup>2</sup> that spans more than one country: all are assessed as Least Concern. Notably, *Valdivia serrata* is found in nine countries (Bolivia, Brazil, Colombia, Ecuador, Peru, Venezuela, Guyana, Suriname, French Guiana), *Poppiana dentata* in eight countries (Brazil, Colombia, French Guiana, Guyana, Nicaragua, Suriname, Trinidad and Tobago, Venezuela), and *Sylviocarcinus pictus* in seven countries (Colombia, Brazil, Bolivia, Peru, Argentina, French Guiana, Guyana). *Dilocarcinus pagei* is found in six countries (Argentina, Brazil, Bolivia, Colombia, Paraguay, Peru), and *Sylviocarcinus devillei* and *Moreirocarcinus emarginatus* are both found in 5 countries (Bolivia, Brazil, Colombia, Ecuador, and Peru; and Brazil, Colombia,

Ecuador, Peru, and Venezuela respectively). Four species are found in four countries: *Trichodactylus kensleyi* (in Argentina, Brazil, Paraguay and Uruguay) and *Poppiana argentiniana*, *Valdivia camerani*, and *Trichodactylus borellianus* (in Argentina, Bolivia, Brazil, and Paraguay). Seven species are found in three countries (*Rotundovaldivia latidens*, *Sylviocarcinus maldonadoensis*, *S. australis*, *Trichodactylus faxoni*, *T. panoplus*, *Valdivia cururuensis* and *Zilchiopsis oronensis*), four species are found in two countries (*Bottiella niceforei*, *B. cucutensis*, *Forsteria venezuelensis*, *T. quinquedentatus*), and three species are found in one country (*Trichodactylus petropolitanus*, *Valdivia novemdentata*, and *Zilchiopsis collastinensis*).

### Distribution patterns and habitat

Freshwater crabs are found in all major habitat types in the Neotropics (Thieme et al. 2005; Abell et al. 2008) including rivers, rapids, swamps, lakes, caves, and mountain streams and species belonging to both families prefer warm-water habitats year round. The Pseudothelphusidae are typically found in small mountain streams above 400 m above sea level, which may in part reflect their ability to breathe air and live a semi-terrestrial existence. Pseudothelphusids reach their highest diversity in the rivers and streams that drain the highland areas of the cordilleras of Colombia and Ecuador. In contrast, trichodactylids are most abundant in the vast network of lowland rivers of South America that make up the Amazon, Orinoco, Paraguay, and Parana basins (Cumberlidge et al. 2009). Species diversity tends to be highest where vegetation cover is densest and water availability is highest, and there are fewer, or no species found in the more arid ecosystems and those with colder water temperatures. The pattern of the dominance of Pseudothelphusidae in Colombia, Venezuela, Ecuador, Peru, Mexico, Central America, and the Caribbean, and their absence in Bolivia, Uruguay, Paraguay, southern Argentina, and Chile probably reflects an origin somewhere in Colombia and Venezuela, from where they spread out north to Central America and Mexico, east to the Guyanas and Lesser Antilles, and south as far as Brazil and Peru (Rodríguez 1982, 1986).

Similarly the dominance of Trichodactylidae in the Amazon, Orinoco, Paraguay, and Parana River basins and their absence from Chile and from aquatic ecosystems at high altitudes in the Andes elsewhere in South America reflects their preference for an aquatic, rather than a semiterrestrial life (Rodríguez 1982, 1986). However, it is harder to explain the presence of trichodactylids in Central America, Mexico, and Trinidad in localities that lie well outside of the major river basins of the South American subcontinent (Rodríguez 1992).

Twenty-four species of neotropical freshwater crabs are found in caves (22 pseudothelphusids and two trichodactylids), and are either fully adapted to a life in caves, or are visitors to caves and have a distribution that also includes surface waters (Table 3). Twelve species of cave-dwelling freshwater crabs are from Mexico, five from Central America (Belize 1, Guatemala 3, and Costa Rica 1), two from the Caribbean (Cuba 1, Puerto Rico, and St. Croix 1), and five from South America (Colombia, 2 and Venezuela 3).

**Table 3.** Species of Neotropical freshwater crabs that are associated with caves, and the country where each species occurs. The family, subfamily/tribe for each species are also shown, together with its IUCN Red List conservation status. LC = Least Concern, EN = Endangered, DD = Data Deficient. For explanation of Red List categories see IUCN (2003). \* = recently described species, not assessed yet for conservation status.

Family	Tribe	Species	Country	Cons. Status
Pseudothelphusidae	Strengerianini	<i>Neostrengeria charalensis</i>	Colombia	LC
Pseudothelphusidae	Strengerianini	<i>Neostrengeria sketi</i>	Colombia	DD
Pseudothelphusidae	Hypolobocerini	<i>Achlidon puntarenas</i>	Costa Rica	DD
Pseudothelphusidae	Epilobocerinae	<i>Epilobocera cubensis</i>	Cuba	LC
Pseudothelphusidae	Hypolobocerini	<i>Phrygiopilus acanthophallus</i>	Guatemala	DD
Pseudothelphusidae	Potamocarcinini	<i>Typhlopseudothelphusa juberthiei</i>	Guatemala	DD
Pseudothelphusidae	Potamocarcinini	<i>Zilchia falcata</i>	Guatemala	DD
Pseudothelphusidae	Potamocarcinini	<i>Typhlopseudothelphusa acanthochela</i>	Belize	DD
Trichodactylidae	Trichodactylinae	<i>Avotrichodactylus constrictus</i>	Mexico	LC
Trichodactylidae	Trichodactylinae	<i>Rodriguezia mensabak</i>	Mexico	DD
Pseudothelphusidae	Hypolobocerini	<i>Phrygiopilus montebelloensis</i>	Mexico	DD
Pseudothelphusidae	Hypolobocerini	<i>Phrygiopilus yoshibensis</i>	Mexico	DD
Pseudothelphusidae	Potamocarcinini	<i>Odontothelphusa monodontis</i>	Mexico	DD
Pseudothelphusidae	Potamocarcinini	<i>Potamocarcinus leptomelus</i>	Mexico	DD
Pseudothelphusidae	Pseudothelphusini	<i>Pseudothelphusa mexicana</i>	Mexico	DD
Pseudothelphusidae	Pseudothelphusini	<i>Tehuana complanata</i>	Mexico	DD
Pseudothelphusidae	Potamocarcinini	<i>Typhlopseudothelphusa hyba</i>	Mexico	DD
Pseudothelphusidae	Potamocarcinini	<i>Typhlopseudothelphusa mocinoi</i>	Mexico	EN
Pseudothelphusidae	Potamocarcinini	<i>Sylvathelphusa cavernicola</i>	Mexico	*
Pseudothelphusidae	Potamocarcinini	<i>Villalobosus lopezformenti</i>	Mexico	DD
Pseudothelphusidae	Epilobocerinae	<i>Epilobocera sinuatifrons</i>	Puerto Rico	LC
Pseudothelphusidae	Strengerianini	<i>Chaceus caecus</i>	Venezuela	DD
Pseudothelphusidae	Strengerianini	<i>Chaceus turikensis</i>	Venezuela	DD
Pseudothelphusidae	Pseudothelphusini	<i>Rodriguezus garmani</i>	Venezuela	LC

Freshwater crab endemism (Table 4)

The majority of freshwater crab species found in Colombia (81%) are country endemics: 78 species of pseudothelphsids and 5 species of trichodactylids (Cumberlidge et al. 2009). The high degree of endemism in Colombia’s freshwater crab fauna at the species level (83 of 102 species, 81%) is also seen to a lesser extent at the genus level (9 of 21 genera, 43%), but not at the family level (neither of the two families are exclusively found in Colombia). Many of the Colombian endemics are found in the isolated mountain streams and in the middle stretches of rivers associated with rainforest (Campos 2005). Mexico’s freshwater crab fauna has the next highest number of country endemics (66 of 67 species, 97%) that is also seen at the genus level (13 of 17 genera, 76%), but not at the family level (neither of the two families are exclusively found in Mexico). Many of the Mexican endemics are found in the isolated mountain streams and in the middle stretches of rivers associated with the rainforests and caves

**Table 4.** Countries that host endemic species of Neotropical freshwater crabs. First number indicates the number of species of Pseudothelphusidae (P), the second number indicates the same data for the Trichodactylidae (T).

Country	No. Endemic Sp. (P, T)
Colombia	78, 5
Mexico	62, 4
Venezuela	22, 0
Panama	12, 1
Ecuador	11, 0
Costa Rica	9, 0
Cuba	9, 0
Guatemala	9, 0
Brazil	7, 10
Peru	6, 3
El Salvador	3, 0
Nicaragua	3, 0
Belize	2, 0
Dominican Republic	2, 0
Guyana	2, 0
Haiti	1, 0
Honduras	1, 0
Suriname	1, 0
Trinidad	1, 0
Bolivia	0, 1

(Villalobos and Alvarez 2010). All species of freshwater crabs found in Cuba and Hispaniola are island endemics.

### Conservation status

The conservation status of the 298 species of freshwater crabs known from the Neotropical region in 2009 was evaluated against the IUCN (2003) Red List criteria (version 3.1) and is available on the Red List site ([www.iucn.redlist.org](http://www.iucn.redlist.org)). These results were summarized by Cumberlidge et al. (2009). The conservation status of species described since that work (Table 1) has not yet been assessed. Unfortunately, about half of all Neotropical freshwater crabs (148 species) are too poorly known to submit to the assessment protocols and are listed as Data Deficient, notably some 80% of the diverse Mexican fauna (Tables 5, 6). Of the remaining 150 species that could be assessed (belonging to 22 genera and 2 families) the majority (107 species) were found to be of Least Concern, and most of these species either live in rivers, marshy lowlands, or in mountain streams in the forested highlands (Cumberlidge et al. 2009). For the Pseudothelphusidae, 111 out of 251 species were assessed (140 species were data deficient), and of these, 72 species of Least Concern, 2 species were Critically Endan-

**Table 5.** Summary of the Red List assessments of the two families of Neotropical freshwater crabs based on data from Cumberlidge et al. (2009). Assessed = number of species that could be assessed using the IUCN Red List protocols; Thr. = total number of species assessed in one of the threatened categories (VU = Vulnerable, EN = Endangered, CR = Critically Endangered); LC = Least Concern, NT = Near Threatened, DD = Data Deficient. For explanation of Red List categories see IUCN (2003).

Family	Total Sp.	Assessed	Thr.	Thr. (%)	VU	EN	CR	LC	NT	DD
Pseudothelphusidae	251	111	38	34.2	33	3	2	72	1	140
Trichodactylidae	47	39	4	10.3	3	1	0	35	0	8
Total	298	150	42	28.3	36	4	2	107	1	148

**Table 6.** Summary of the countries in the Neotropical region that have threatened species of freshwater crabs based on data from Cumberlidge et al. (2009). Assessed = number of species that could be assessed using the IUCN Red List protocols; Thr. = total number of species assessed in one of the threatened categories (VU = Vulnerable, EN = Endangered, CR = Critically Endangered); LC = Least Concern, NT = Near Threatened, DD = Data Deficient.

Country	Total Sp.	Assessed	Thr.	Thr. (%)	VU	EN	CR	LC	NT	DD
Colombia	101	60	14	23.3	13	0	1	45	1	41
Venezuela	42	31	7	22.6	7	0	0	24	0	11
Mexico	63	19	6	31.6	2	3	1	13	0	44
Ecuador	27	19	5	36.8	5	0	0	12	0	8
Peru	25	21	3	14.3	3	0	0	0	0	4
El Salvador	8	6	2	33.3	2	0	0	4	0	2
Costa Rica	13	7	2	28.6	2	0	0	5	0	6
Brazil	45	38	2	10.5	1	1	0	34	0	7
Suriname	11	10	1	10.0	1	0	0	0	0	1
Honduras	6	6	1	33.3	1	0	0	4	0	0
Haiti	1	1	1	100.0	1	0	0	0	0	0
Guatemala	13	7	1	28.6	1	0	0	5	0	6
Dominican Rep.	2	2	1	100.0	1	0	0	0	0	0
Cuba	9	9	1	11.0	1	0	0	0	0	0

gered, 3 species were Endangered Critically Endangered, 3 species were Endangered, and 33 species were Vulnerable (Tables 5, 6). For the Trichodactylidae, 39 out of 47 species were assessed (8 species were Data Deficient), and of these, 35 species were of Least Concern, one species was Endangered, and three were Vulnerable (Table 6). For both families combined, 42 of the 150 assessed species of Neotropical freshwater crabs (28%) were listed in one of three threatened categories, either as Critically Endangered (2 species), Endangered (4 species) or Vulnerable (36 species) (Table 7). One species was assessed as near threatened, and no species of Neotropical freshwater crabs was confirmed as Extinct or Extinct in the Wild. However, it should be noted that a species cannot be formally assessed as extinct until exhaustive surveys probing its disappearance have been carried out. Because conservation status could not be assigned to large numbers of Data Deficient species (148), the proportion of threatened species in Cumberlidge

**Table 7.** Threatened species of Neotropical freshwater crabs. P = Pseudothelphusidae, T = Trichodactylidae. For explanation of Red list categories see IUCN (2003).

Family/Subfamily/Tribe	Taxon	IUCN Red List Category	Distribution
T. (Trichodactylinae)	<i>Trichodactylus crassus</i>	EN B1ab(iii)	Brazil
P. (Kingsleyini)	<i>Kingsleya siolii</i>	VU B1ab(iii); D2	Brazil, Suriname
P. (Kingsleyini)	<i>Fredius platyacanthus</i>	VU B1ab(iii); D2	Brazil, Venezuela
P. (Kingsleyini)	<i>Fredius stenolobus</i>	VU B1ab(iii); D2	Brazil, Venezuela
P. (Hypolobocerini)	<i>Hypolobocera alata</i>	VU B1ab(iii); D2	Colombia
P. (Hypolobocerini)	<i>Hypolobocera andagoensis</i>	VU B1ab(iii); D2	Colombia
P. (Hypolobocerini)	<i>Hypolobocera barbacensis</i>	VU B1ab(iii); D2	Colombia
P. (Hypolobocerini)	<i>Hypolobocera cajambrensis</i>	VU B1ab(iii); D2	Colombia
P. (Hypolobocerini)	<i>Hypolobocera rotundilobata</i>	VU B1ab(iii); D2	Colombia
P. (Hypolobocerini)	<i>Hypolobocera velezi</i>	VU B1ab(iii); D2	Colombia
P. (Hypolobocerini)	<i>Moritschus altaquerensis</i>	VU B1ab(iii); D2	Colombia
P. (Kingsleyini)	<i>Fredius granulatus</i>	VU B1ab(iii); D2	Colombia
P. (Strengerianini)	<i>Chaceus ibiricensis</i>	VU B1ab(iii)+2ab(iii); D2	Colombia
P. (Strengerianini)	<i>Strengeriana antioquensis</i>	CR B1ab(i,ii,iii,iv,v)+2ab(i,ii,iii,iv,v)	Colombia
T. (Dilocarcininae)	<i>Bottiella medemi</i>	VU B1ab(iii)+2ab(iii)	Colombia
P. (Hypolobocerini)	<i>Hypolobocera exuca</i>	VU B1ab(iii); D2	Colombia, Ecuador
P. (Hypolobocerini)	<i>Lindacatalina sumacensis</i>	VU B1ab(iii); D2	Colombia, Ecuador
T. (Dilocarcininae)	<i>Bottiella cucutensis</i>	VU B1ab(iii)+2ab(iii)	Colombia, Venezuela
P. (Potamocarcini)	<i>Allacanthos pittieri</i>	VU B1ab(iii); D2	Costa Rica
P. (Potamocarcini)	<i>Ptychophallus tristani</i>	VU B1ab(iii)+2ab(iii); D2	Costa Rica
P. (Epilobocerinae)	<i>Epilobocera wetherbeeii</i>	VU B1ab(iii); D2	Dominican Republic
P. (Hypolobocerini)	<i>Hypolobocera delsolari</i>	VU B1ab(iii); D2	Ecuador
P. (Hypolobocerini)	<i>Hypolobocera rathbuni</i>	VU B1ab(iii); D2	Ecuador
P. (Hypolobocerini)	<i>Moritschus ecuadorensis</i>	VU B1ab(iii); D2	Ecuador
P. (Hypolobocerini)	<i>Elsalvadoria zurstrasseni</i>	VU B1ab(iii)+2ab(iii); D2	El Salvador
P. (Hypolobocerini)	<i>Raddaus mertensi</i>	VU D2	El Salvador, Guatemala, Honduras
P. (Hypolobocerini)	<i>Phrygiopilus acanthophallus</i>	VU B1ab(iii); D2	Guatemala
P. (Epilobocerinae)	<i>Epilobocera haytensis</i>	VU B1ab(iii)	Haiti, Dominican Rep.
P. (Potamocarcini)	<i>Potamocarcinus roatensis</i>	VU D2	Honduras
P. (Potamocarcini)	<i>Potamocarcinus hartmanni</i>	VU B1ab(ii)+2ab(ii); D2	Mexico
P. (Potamocarcini)	<i>Typhlopseudothelphusa mocinoi</i>	EN B1ab(iii)+2ab(iii)	Mexico
P. (Pseudothelphusini)	<i>Tehuana lamothiei</i>	EN B1ab(ii,iii)+2ab(ii,iii)	Mexico
P. (Pseudothelphusini)	<i>Tehuana poglayenorum</i>	EN B1ab(ii,iii)+2ab(ii,iii)	Mexico
P. (Pseudothelphusini)	<i>Tehuana veracruzana</i>	CR B1ab(i,ii,iii,iv,v)+2ab(i,ii,iii,iv,v)	Mexico
T. (Trichodactylinae)	<i>Avotrichodactylus oaxensis</i>	VU B1ab(iii)+2ab(iii)	Mexico
P. (Hypolobocerini)	<i>Hypolobocera gracilignatha</i>	VU D2	Peru
P. (Hypolobocerini)	<i>Hypolobocera lamercedes</i>	VU B1ab(iii); D2	Peru
P. (Hypolobocerini)	<i>Hypolobocera peruviana</i>	VU D2	Peru
P. (Kingsleyini)	<i>Microthelphusa forcarti</i>	VU B1ab(iii); D2	Venezuela
P. (Kingsleyini)	<i>Neopseudothelphusa fossor</i>	VU B1ab(iii); D2	Venezuela
P. (Kingsleyini)	<i>Rodriguezus trujillensis</i>	VU B1ab(iii)+2ab(iii); D2	Venezuela

et al. (2009) (32%) is almost certainly an underestimation. This means that the number of threatened species could well increase should a Data Deficient species subsequently be assessed as threatened (Cumberlidge et al. 2009). Despite the large numbers of specimens available very little information is available on population levels and trends, except for a qualitative estimate (e.g., whether common or rare) based on the number of sites at which a species is present and its relative abundance at each site.

With only 42 out of 298 species of Neotropical freshwater crabs known in 2009 assessed as threatened with global extinction (Tables 5–7), the region's largely endemic freshwater crab fauna (Table 4) does not appear at first sight to be in immediate trouble compared with other aquatic groups found in the same freshwater habitats (e.g., fish, molluscs, and dragonflies) (Collen et al. 2008). The 107 species of Neotropical freshwater crabs judged to be of Least Concern (35 species of trichodactylids, and 72 species of pseudothelphusids) have a wide distribution in rivers, lakes, highland streams and lowland wetlands, and appear to be tolerant of changes in land use that affect aquatic ecosystems.

Threats to Neotropical freshwater crabs include habitat destruction driven by increasing agriculture, the demands of increasing industrial development, and the alteration of fast flowing rivers for the creation of hydroelectric power (Cumberlidge et al. 2009). Even species assessed as Least Concern could suffer a catastrophic decline should there be abrupt changes in land development, hydrology, or pesticide-use regimes. Species with a narrow distribution are vulnerable to extreme population fragmentation and could suffer a rapid decline and even extinction in a relatively short time should dramatic changes in land-use affect their habitat. The 148 species of Neotropical freshwater crabs judged to be Data Deficient are mostly rare species, and their conservation status needs to be re-evaluated once more information comes to light. It is hoped that prioritizing species for conservation action through the Red List assessment process will lead to the development of conservation recovery plans for threatened species in the future.

## Conclusions

This study confirms that freshwater crabs are completely absent from all parts of the Nearctic region (even in the neighboring warmer parts of northern Mexico and southern Florida, USA), and that the northern limits of their distribution coincide with the Nearctic/Neotropical boundary in Mexico and the presence of arid regions lacking permanent freshwaters (i.e., the Sonoran and Chihuahuan Deserts). Species richness of freshwater crabs increases south of this boundary in Mexico and reaches a peak in the States of Veracruz, Tabasco, Oaxaca, and Chiapas. The uneven pattern of species-richness – high in Mexico, low in Central America, and high in Colombia – is difficult to explain by simply invoking an even spread of species out of a center of origin somewhere in Colombia–Venezuela (Rodríguez 1982, 1992), and further studies are necessary. The relatively low species richness of freshwater crabs in Brazil is unexpected given its vast area and rich biodiversity, and may reflect the dominance of trichodactylids and a general paucity of species-rich pseudothelphusids in the Brazilian fauna. The abundance of freshwater crabs

in Colombia, Ecuador, and Peru coincides with a freshwater habitat species diversity hotspot (that is defined by overlapping distributions of stenotopic species based on combined data from vertebrates and invertebrates (Collen et al. 2008)). All countries in the Neotropical region are still in a phase of exploration and steep rises in species numbers of pseudothelphusids are expected once more surveys have been completed.

The Neotropics are the second most diverse region in the world for freshwater crabs, and there are 42 species from 14 countries within this region that may be threatened with extinction. The finding that 34% of pseudothelphusids and 10% of trichodactylids are threatened, and that many others are poorly known, can be used to develop a conservation strategy for these species, and supply a focus for future research efforts (Cumberlidge et al. 2009). Although some species of Neotropical freshwater crabs have been quite well studied a clear majority are known only from either the type locality or from just a few localities, and in these cases further collections are necessary to ascertain their actual distributions. The restricted range of many species from the Neotropics, together with the on-going human induced loss of habitat in many parts of the region, are a cause for concern for the long-term security of elements of this fauna. Conservation activities should therefore be aimed primarily at preserving the integrity of sites and habitats while at the same time closely monitoring key populations. Significant areas of the Neotropics still remain insufficiently explored, and the focus of new efforts should be on new species discovery through increased collection in remote areas and the refinement of taxonomic skills. These efforts should also focus on conserving threatened species, and on seeking out data deficient species. Gathering current data on distribution, natural history, population trends, threats, and endemism of the Neotropical region's highly diverse freshwater crabs will enable the updating of the IUCN Red Lists, and the development of conservation strategies for this understudied diverse and potentially highly threatened fauna.

## Acknowledgements

The expert contributions of Célio Magalhães, Martha Campos and the second and third authors to the global assessment for the Neotropical freshwater crabs are recognized here. Amy Peterson (Northern Michigan University, USA) is thanked for her assistance with the georeferencing of large numbers of pseudothelphusid and trichodactylid localities. Gema Armendáriz and Benjamin Quiroz (Instituto de Biología, UNAM, Mexico) are thanked for adding records to the freshwater crab database and for creating the maps.

## References

- Abell R, Thieme ML, Revenga C, Bryer M, Kottelat M, Bogutskaya N, Coad B, Mandrak N, Balderas SC, Bussing W, Stiassny MLJ, Skelton P, Allen GR, Unmack P, Naseka A, Ng

- R, Sindorf N, Robertson J, Armijo E, Higgins JV, Heibel TJ, Wikramanayake E, Olson D, Lopez HL, Reis RE, Lundberg JG, Perez MHS, Petry P (2008) Freshwater ecoregions of the world: a new map of biogeographic units for freshwater biodiversity conservation. *BioScience* 58(5): 403–414. doi: 10.1641/B580507
- Acha PN, Szyfres B (2001) Zoonoses and Communicable Diseases Common to Man and Animals. Pan American Health Organization, 3<sup>rd</sup> edition. Washington, DC, 384 pp.
- Alvarez F (1987) *Pseudothelphusa mexicana*, a new freshwater crab from the State of Guerrero, Mexico (Brachyura: Pseudothelphusidae). *Proceedings of the Biological Society of Washington* 100: 1–4.
- Alvarez F (1989) *Smalleyus tricristatus*, new genus, new species, and *Pseudothelphusa parabeliana*, new species (Brachyura: Pseudothelphusidae) from Los Tuxtlas, Veracruz, Mexico. *Proceedings of the Biological Society of Washington* 102: 45–49.
- Alvarez F, Villalobos JL (1990) *Pseudothelphusa galloi*, new species of freshwater crab (Brachyura: Pseudothelphusidae) from southwestern Mexico. *Proceedings of the Biological Society of Washington* 103: 103–105.
- Alvarez F, Villalobos JL (1991) A new genus and two new species of freshwater crabs from Mexico, *Odonthelphusa toninae* and *Stygothelphusa lopezformenti* (Crustacea: Brachyura: Pseudothelphusidae). *Proceedings of the Biological Society of Washington* 104: 288–294.
- Alvarez F, Villalobos JL (1994) Two new species and one new combination of freshwater crabs from Mexico (Crustacea: Brachyura: Pseudothelphusidae). *Proceedings of the Biological Society of Washington* 107: 729–737.
- Alvarez F, Villalobos JL (1995) Lista anotada de los Tipos de la Colección de Crustáceos del Instituto de Biología, Universidad Nacional Autónoma de México. 1. Cangrejos Verdaderos del Infraorden Brachyura. *Revista de la Sociedad Mexicana de Historia Natural* 46: 87–96.
- Alvarez F, Villalobos JL (1996) Especie nueva de cangrejo de agua dulce del género *Pseudothelphusa* (Brachyura: Pseudothelphusidae) de Guerrero, México. *Anales del Instituto de Biología, Universidad Nacional Autónoma de México* 67: 297–302.
- Alvarez F, Villalobos JL (1997a) *Pseudothelphusa ayutlaensis*, a new species of freshwater crab (Brachyura: Pseudothelphusidae) from Mexico. *Proceedings of the Biological Society of Washington* 110: 388–392.
- Alvarez F, Villalobos JL (1997b) Pseudothelphusidae y Trichodactylidae (cangrejos). In: González-Soriano E, Dirzo R, Vogt RC (Eds) *Historia Natural de la Región de Los Tuxtlas, Veracruz*. Primera Edición, UNAM-CONABIO, 415–418.
- Alvarez F, Villalobos JL (1998) Six new species of freshwater crabs (Brachyura: Pseudothelphusidae) from Chiapas, Mexico. *Journal of Crustacean Biology* 18: 187–198. doi: 10.2307/1549533
- Alvarez F, Villalobos JL, Lira E (1996) Decapoda. In: Llorente-Bousquets J, García-Aldrete AN, González-Soriano E (Eds) *Biodiversidad, Taxonomía y Biogeografía de Artrópodos de México: Hacia Una Síntesis de su Conocimiento*. Primera edición. Universidad Nacional Autónoma de México, México, D.F., 103–129.
- Alvarez F, Villalobos JL, Moreno E (2012a) *Pseudothelphusa zongolicae* (Decapoda, Pseudothelphusidae), a new species of freshwater crab from Veracruz, Mexico. *Crustaceana* 85: 1541–1547. doi: 10.1163/15685403-00003132

- Alvarez F, Villalobos JL, Rojas Y, Robles R (1999) Lista y comentarios sobre los crustáceos decápodos de Veracruz, México. *Anales del Instituto de Biología, Universidad Nacional Autónoma de México* 70: 1–27.
- Alvarez F, Villalobos JL, Armendáriz G, Hernández C (2012b) Relación biogeográfica entre cangrejos dulceacuícolas y acociles a lo largo de la mexicana de transición: revaluación de la hipótesis de Rodríguez (1986). *Revista Mexicana de Biodiversidad* 83: 1073–1083.
- Bott R (1967) Fluss-Krabben aus dem westlichen Mittelamerika (Crust. Decap.). *Senckenbergiana Biologica* 48: 373–380.
- Bott R (1968) Fluss-Krabben aus dem östlichen Mittel-Amerika und von den Grossen Antillen (Crustacea, Decapoda). *Senckenbergiana Biologica* 49: 39–49.
- Bott R (1969) Preadaption, Evolution und Besiedlungsgeschichte der Süßwasserkrabben der Erde. *Natur und Museum* 99: 266–275.
- Bott R (1970) Betrachtungen über die Entwicklungsgeschichte und Verbreitung der Süßwasserkrabben nach der Sammlung des Naturhistorischen Museums in Genf/Schweiz. *Review Suisse de Zoologie* 77: 327–344.
- Campos MR (1989) Nuevas especies de cangrejos de agua dulce del genero *Hypolobocera* (Crustacea: Decapoda: Pseudothelphusidae) para Colombia. *Trianea* 3: 143–147.
- Campos MR (1992) New species of fresh-water crabs of the genus *Neostrengeria* (Crustacea: Decapoda: Pseudothelphusidae) from Colombia. *Proceedings of the Biological Society of Washington* 105(3): 540–554.
- Campos MR (1995) A new species of freshwater crab of the genus *Strengeriana* from Colombia (Crustacea: Decapoda: Pseudothelphusidae). *Proceedings of the Biological Society of Washington* 108(1): 98–101.
- Campos MR (1997) Primeros registros de la tribu Kingsleyini (Crustacea: Decapoda: Pseudothelphusidae) para Colombia. *Caldasia* 19(1–2): 269–277.
- Campos MR (1999) A new species of freshwater crab of the genus *Strengeriana* Pretzmann, 1971, from Colombia (Crustacea: Decapoda: Pseudothelphusidae). *Proceedings of the Biological Society of Washington* 112(2): 405–409.
- Campos MR (2000) *Neostrengeria binderi*, a new species of pseudothelphusid crab from the eastern Andes of Colombia (Crustacea: Decapoda: Brachyura). *Proceedings of the Biological Society of Washington* 113(2): 401–405.
- Campos MR (2001) A new genus and species of freshwater crab from Colombia (Crustacea: Decapoda: Pseudothelphusidae). *Proceedings of the Biological Society of Washington* 114(4): 938–943.
- Campos MR (2002) *Dilocarcinus laevifrons* Moreira, 1901: a complete description (Decapoda; Brachyura: Trichodactylidae). *Revista de la Academia Colombiana de Ciencias Exactas, Físicas y Naturales* 26(99): 285–288.
- Campos MR (2003a) A new species of freshwater crab of the genus *Potamocarcinus* H. Milne Edwards, 1853 (Crustacea: Brachyura: Pseudothelphusidae) from Colombia. *Revista de la Academia Colombiana de Ciencias Exactas, Físicas y Naturales* 27(103): 283–286.
- Campos MR (2003b) A review of the freshwater crabs of the genus *Hypolobocera* Ortmann, 1897 (Crustacea: Decapoda: Brachyura: Pseudothelphusidae), from Colombia. *Proceedings of the Biological Society of Washington* 116(3): 754–802.

- Campos MR (2004) *Neostrengeria lemaitrei*, a new species of freshwater crab from Colombia (Crustacea: Decapoda: Pseudothelphusidae), and the vertical distribution of the genus. *Proceedings of the Biological Society of Washington* 117(3): 363–367.
- Campos MR (2005) Freshwater crabs from Colombia: A taxonomic and distributional study. Academia Colombiana de Ciencias Exactas, Físicas y Naturales. Colección Jorge Alvarez Lleras 24: 1–363.
- Campos MR (2010a) Estudio taxonómico de los crustáceos decápodos de agua dulce (Trichodactylidae, Pseudothelphusidae) de Casanare, Colombia. *Revista de la Academia Colombiana de Ciencias* 34: 257–266.
- Campos MR (2010b) A new species of freshwater crab of the genus *Neostrengeria* Pretzmann, 1965, from Colombia (Crustacea: Decapoda: Pseudothelphusidae). *Zootaxa* 2482: 64–68.
- Campos MR (2011) A new species of freshwater crab of the genus *Neostrengeria* Pretzmann, 1965, from Meta Department, Colombia (Crustacea: Decapoda: Pseudothelphusidae). *Caldasia* 33: 659–664.
- Campos MR, Lemaitre R (1998) A new freshwater crab of the genus *Neostrengeria* Pretzmann, 1965, from Colombia (Crustacea: Decapoda: Brachyura: Pseudothelphusidae), with a key to the species of the genus. *Proceedings of the Biological Society of Washington* 111: 899–907.
- Campos MR, Lemaitre R (2002) A new species of freshwater crab of the genus *Potamocarcinus* H. Milne-Edwards, 1853 (Crustacea: Decapoda: Brachyura: Pseudothelphusidae) from Panama. *Proceedings of the Biological Society of Washington* 115: 600–604.
- Campos MR, Pedraza M (2006) A new species of freshwater crab of the genus *Strengeriana* Pretzmann, 1971, from Colombia (Crustacea: Decapoda: Pseudothelphusidae), with an updated key to the species of the genus. *Proceedings of the Biological Society of Washington* 119: 557–562. doi: 10.2988/0006-324X(2006)119[557:ANSOFC]2.0.CO;2
- Campos MR, Pedraza M (2008) Two new species of freshwater crab of the genus *Neostrengeria* Pretzmann, 1965, from Colombia (Crustacea: Decapoda: Pseudothelphusidae), with an updated key of the species of the genus. *Caldasia* 30: 457–468
- Campos MR, Rodríguez G (1984) New species of freshwater crabs (Crustacea: Decapoda: Pseudothelphusidae) from Colombia. *Proceedings of the Biological Society of Washington* 97(3): 538–543.
- Campos MR, Rodríguez G (1985) A new species of *Neostrengeria* (Crustacea: Decapoda: Pseudothelphusidae) with notes on geographical distribution of the genus. *Proceedings of the Biological Society of Washington* 98(3): 718–727.
- Campos MR, Rodríguez G (1988) Notes on the freshwater crabs of the genus *Moritschus* Pretzmann, 1965 (Crustacea: Decapoda: Pseudothelphusidae) with description of *M. narinnensis* from southern Colombia. *Proceedings of the Biological Society of Washington* 101(3): 640–643.
- Campos MR, Rodríguez G (1993) Three new species of *Strengeriana* from Colombia (Crustacea: Decapoda: Pseudothelphusidae). *Proceedings of the Biological Society of Washington* 106(3): 508–513.
- Campos MR, Rodríguez G (1995) Two new species of freshwater crabs of the genus *Hypolobocera* from Colombia (Crustacea: Decapoda: Pseudothelphusidae). *Proceedings of the Biological Society of Washington* 108(4): 649–655.

- Campos MR, Valencia (2004) Two new species of freshwater crabs of the genus *Chaceus* Pretzmann, 1965 from the Serrania De Perija of Colombia (Crustacea: Decapoda: Pseudothelphusidae). *Proceedings of the Biological Society of Washington* 117: 35–41.
- Cardona L, Campos M (2012) A new species of freshwater crab of the genus *Phallangothelphusa* Pretzmann, 1965, from Colombia (Crustacea: Decapoda: Pseudothelphusidae). *Zootaxa* 3515: 83–88.
- Collen B, Ram M, Dewhurst N, Clausnitzer V, Kalkman V, Cumberlidge N, Baillie JEM (2008) Broadening the coverage of biodiversity assessments. In: Viè J-C, Hilton-Taylor C, Stuart SN (Eds) *The 2008 Review of the IUCN Red List of Threatened Species*. IUCN Gland, Switzerland, 1–12.
- Cumberlidge N (2007) A new species of freshwater crab of the genus *Microthelphusa* (Brachyura: Pseudothelphusidae) from a remote isolated cloud forest on a tabletop mountain in western Guyana, South America. *Zootaxa* 1447: 57–62.
- Cumberlidge N, Ng PKL, Yeo DCJ, Magalhães C, Campos MR, Alvarez F, Naruse T, Daniels SR, Esser LJ, Collen B, Attipoe FYK, Ba FJC, Darwall W, McIvor A, Ram M (2009) Freshwater crabs and the biodiversity crisis: importance, threats, status, and conservation challenges. *Biological Conservation* 142: 1665–1673. doi: 10.1016/j.biocon.2009.02.038
- Herbst JFW (1783) *Versuch einer Naturgeschichte der Krabben und Krebse, nebst einer systematischen Beschreibung ihrer verschiedenen Arten*. Gottlieb August Lange, Berlin and Stralsund, 1–3: 1–515.
- Hobbs HH Jr. (1980) A new pseudothelphusid crab from the state of Jalisco, Mexico. *Proceedings of the Biological Society of Washington* 93: 357–361.
- Hobbs HH III (1991) A new pseudothelphusid crab from a cave in southern Costa Rica (Decapoda: Brachyura). *Proceedings of the Biological Society of Washington* 104: 295–298.
- Huidobro L, Morrone JJ, Villalobos JL, Alvarez F (2006) Distributional patterns of freshwater taxa (fishes, crustaceans and plants) from the Mexican transition zone. *Journal of Biogeography* 33: 731–741. doi: 10.1111/j.1365-2699.2005.01400.x
- IUCN (2003) *Guidelines for Application of IUCN red list Criteria at Regional Levels: Version 3.0*. IUCN Species Survival Commission. IUCN, Gland, Switzerland and Cambridge, UK.
- Magalhães C (1986) Revisão taxonômica dos caranguejos de água doce brasileiros da família Pseudothelphusidae (Crustacea: Decapoda). *Amazoniana* 9(4): 609–636.
- Magalhães C (2003) The occurrence of freshwater crabs (Crustacea: Decapoda: Pseudothelphusidae) in the Rio Xingu, Amazon Region, Brazil, with description of a new species of Pseudothelphusidae. *Amazoniana* 17(3,4): 377–386.
- Magalhães C (2010) A new species of freshwater crab of the genus *Microthelphusa* (Decapoda, Pseudothelphusidae) from a tepui in the Serra Do Aracá, state of Amazonas, Brazil. In: Fransen C, de Grave S, Ng P (Eds) *Studies on Malacostraca: Lipke Bijdeley Holthuis Memorial Volume*. *Crustaceana Monographs* 14: 453–460.
- Magalhães C, Türkay M (1986) *Brasiliiothephusa*, a new Brazilian freshwater-crab genus (Crustacea: Decapoda: Pseudothelphusidae). *Senckenbergiana biologica* 66(4/6): 371–376.
- Magalhães C, Türkay M (1996a) Taxonomy of the neotropical freshwater crab family Trichodactylidae. I. The generic system with description of some new genera (Crustacea: Decapoda: Brachyura). *Senckenbergiana Biologica* 75(1/2): 63–95.

- Magalhães C, Türkay M (1996b) Taxonomy of the neotropical freshwater crab family Trichodactylidae. II. The genera *Forsteria*, *Melocarcinus*, *Sylviocarcinus*, and *Zilchiopsis* (Crustacea: Decapoda: Brachyura). *Senckenbergiana Biologica* 75(1/2): 97–130.
- Magalhães C, Türkay M (1996c) Taxonomy of the neotropical freshwater crab family Trichodactylidae. III. The genera *Fredilocarcinus*, and *Goyazana* (Crustacea: Decapoda: Brachyura). *Senckenbergiana Biologica* 75(1/2): 131–142.
- Magalhães C, Türkay M (2008a) Taxonomy of the neotropical freshwater crab family Trichodactylidae. IV. The genera *Dilocarcinus* and *Poppiana* (Crustacea: Decapoda: Trichodactylidae). *Senckenbergiana Biologica* 88(2): 185–215.
- Magalhães C, Türkay M (2008b) Taxonomy of the neotropical freshwater crab family Trichodactylidae. V. The genera *Bottiella* and *Rotundovaldivia* (Crustacea: Decapoda: Trichodactylidae). *Senckenbergiana biologica* 88(2): 217–230.
- Magalhães C, Türkay M (2008c) A new species of *Kingsleya* from the Yanomami Indians area in the Upper Rio Orinoco, Venezuela. (Crustacea: Decapoda: Brachyura: Pseudothelphusidae). *Senckenbergiana biologica* 88(2): 1–7.
- Magalhães C, Türkay M (2010) A new freshwater crab of the genus *Brasiliothelphusa* Magalhães and Türkay, 1986 from Rio Aripuaná, southern Amazon Region, Brazil (Decapoda: Pseudothelphusidae). *Nauplius* 18(2): 103–108.
- Magalhães C, Türkay M (2012) Taxonomy of the Neotropical freshwater crab family Trichodactylidae, VI. The genera *Avotrichodactylus* and *Rodriguezia* (Crustacea: Decapoda: Trichodactylidae). *Nauplius* 20(1): 27–40. doi: 10.1590/S0104-64972012000100004
- Magalhães C, Campos MR, Türkay M (2013) Freshwater crabs from eastern Panamá: a new species of *Potamocarcinus* H. Milne Edwards, 1853, and new records of two little-known species (Crustacea: Decapoda: Pseudothelphusidae, Trichodactylidae). *Zootaxa* 3702(4): 348–356. doi: 10.11646/zootaxa.3702.4.2
- Magalhães C, Lara LR, Wehrtmann IS (2010) A new species of freshwater crab of the genus *Alacanthos* Smalley, 1964 (Crustacea: Decapoda: Pseudothelphusidae) from southern Costa Rica, Central America. *Zootaxa* 2604: 52–60.
- Ng PKL, Guinot D, Davie PJF (2008) An annotated checklist of extant brachyuran crabs of the world. *Systema Brachyurorum*, Part I. *Raffles Bulletin of Zoology (Supplement)* 17: 1–286.
- Ojeda JC, Alvarez F, Villalobos JL (2013) Revisión de los cangrejos de agua dulce de la familia Trichodactylidae de México. In: López M, Mejía LM (Eds) *La Carcinología en México: El Legado del Dr. Alejandro Villalobos 30 Años Después*. Universidad de Quintana Roo, México, 23–36.
- Ortmann AE (1893) Die Decapoden-Krebse des Strasbourg Museums, mit besonderer Berücksichtigung der von Herrn Dr. Duderlein bei Japan und bei den Liu-Kiu-Inseln gesammelten und zur Zeit im Strassburger Museum aufbewahrten Formen. VII. Theil. Abtheilung: Brachyura (Brachyura genuina Boas) II. Unterabtheilung: Cancroidea, 2. Section: Cancrinea, 1. Gruppe: Cyclometopa. *Zoologische Jahrbücher Abtheilung für Systematik, Geographie und Biologie der Thiere* 7: 411–495.
- Ortmann AE (1897) Carcinologische Studien. *Zoologische Jahrbücher Abtheilung für Systematik, Geographie und Biologie der Thiere* 10: 258–372.
- Pretzmann G (1972) Die Pseudothelphusidae (Crustacea Brachyura). *Zoologica* 42: 1–182.

- Pretzmann G (1975) Zur Systematik der Pseudothelphusidae (Decapoda, Brachyura). *Crustaceana* 27: 294–302. doi: 10.1163/156854074X00820
- Rathbun MJ (1893) Descriptions of new species of American freshwater crabs. *Proceedings of the United States National Museum* 16: 649–661. doi: 10.5479/si.00963801.16-959.649
- Rathbun MJ (1898) A contribution to a knowledge of the fresh-water crabs of America. The Pseudothelphusinae. *Proceedings of the United States National Museum* 21: 507–537. doi: 10.5479/si.00963801.21-1158.507
- Rathbun MJ (1905) Les Crabes d'eau douce (Potamonidae). *Nouvelles Archives du Museum d'Histoire naturelle, Paris, series 4, 7*: 159–321.
- Rodríguez G (1980) Description préliminaire de quelques espèces et genres nouveaux de Crabes d'eau douce de l'Amerique tropicale (Crustacea: Decapoda: Pseudothelphusidae). *Bulletin du Museum Nationale d'Histoire Naturelle, Paris 4<sup>a</sup> séries 2, section A(3)*: 889–894.
- Rodríguez G (1982) Les crabes d'eau douce d'Amerique. Famille des Pseudothelphusidae. *Faune Tropicale* 22: 1–223.
- Rodríguez G (1986) Center of radiation of freshwater crabs in the Neotropics. In: Gore RH, Heck KL (Eds) *Crustacean Biogeography. Crustacean Issues 4*. Balkema, Rotterdam, 51–67.
- Rodríguez G (1992) The freshwater crabs of America. *Faune Tropicale* 31: 1–189.
- Rodríguez G, Campos MR (1989) Cladistic relationships of freshwater crabs of the tribe Stengerianini (Decapoda: Pseudothelphusidae) from the northern Andes, with comments on their biogeography and descriptions of new species. *Journal of Crustacean Biology* 9: 141–156. doi: 10.2307/1548455
- Rodríguez G, Campos MR (1998) A cladistic revision of the genus *Fredius* (Crustacea; Decapoda; Pseudothelphusidae) and its significance to the biogeography of the Guianan lowlands of South America. *Journal of Natural History* 32: 763–765. doi: 10.1080/00222939800770391
- Rodríguez G, Magalhães C (2005) Recent advances in the biology of the Neotropical freshwater crab family Pseudothelphusidae (Crustacea, Decapoda, Brachyura). *Revista Brasileira de Zoologia* 22: 354–365. doi: 10.1590/S0101-81752005000200009
- Rodríguez G, Pereira G (1992) New species, cladistic relationships, and biogeography of the genus *Fredius* (Decapoda: Brachyura: Pseudothelphusidae) from South America. *Journal of Crustacean Biology* 12: 298–311. doi: 10.2307/1549082
- Rodríguez G, Smalley AE (1969) Los cangrejos de agua dulce de México de la familia Pseudothelphusidae (Crustacea, Brachyura). *Anales del Instituto de Biología, Universidad Nacional Autónoma de México* 40: 69–112.
- Rodríguez G, Sternberg R (1998) A revision of the freshwater crabs of the family Pseudothelphusidae (Decapoda: Brachyura) from Ecuador. *Proceedings of the Biological Society of Washington* 111: 110–139.
- Rodríguez G, Suárez H (2004) A revision of the freshwater crabs of the family Pseudothelphusidae (Decapoda: Brachyura) from Peru with notes on the southern limits of the family. *Amazoniana* 18: 11–28.
- Rodríguez G, Campos MR, López B (2002) New species and new records of pseudothelphusid crabs (Crustacea: Brachyura) from Colombia. *Tulane Studies in Zoology and Botany* 31(2): 1–17.

- Smalley AE (1964a) A terminology for the gonopods of the American river crabs. *Systematic Zoology* 13: 28–31. doi: 10.2307/2411434
- Smalley AE (1964b) The river crabs of Costa Rica, and the subfamilies of the Pseudothelphusidae. *Tulane Studies in Zoology* 12: 5–13.
- Smalley AE (1965) Two new freshwater crabs from Nicaragua. *Annals and Magazine of Natural History* 7: 651–656. doi: 10.1080/00222936408651511
- Smalley AE (1970) A new genus of freshwater crabs from Guatemala, with a key to the middle American genera (Crustacea, Decapoda, Pseudothelphusidae). *American Midland Naturalist* 83: 96–106. doi: 10.2307/2424009
- Smalley AE, Rodríguez G (1972) Trichodactylidae from Venezuela, Colombia, and Ecuador (Crustacea: Brachyura). *Tulane Studies in Zoology and Botany* 17: 41–55.
- Smalley AE, Adkison DL (1984) *Disparithelphusa pecki*, a new genus and species of freshwater crab from Mexico (Brachyura: Pseudothelphusidae). *Journal of Crustacean Biology* 4: 127–133. doi: 10.2307/1547901
- Sternberg R, Cumberlidge N (2001) Notes on the position of true freshwater crabs within the brachyrynchan Eubrachyura (Crustacea: Decapoda: Brachyura). *Hydrobiologia* 449: 21–39. doi: 10.1023/A:1017560113027
- Sternberg R, Cumberlidge N, Rodríguez G (1999) On the marine sister groups of the freshwater crabs (Crustacea: Decapoda: Brachyura). *Journal of Zoology, Systematics and Evolution Research* 37: 19–38. doi: 10.1046/j.1439-0469.1999.95092.x
- Thieme ML, Abell RA, Stiassny MJL, Skelton PA, Lehner B (2005) *Freshwater Ecoregions of Africa and Madagascar: A Conservation Assessment*. Island Press, Washington, DC, USA.
- Villalobos JL (2005) *Sistemática de los cangrejos de agua dulce de México, Tribu Pseudothelphusini Ortmann, 1897 (Crustacea: Decapoda: Brachyura: Pseudothelphusidae). Análisis filogenético, biogeográfico y descripción de especies nuevas*. Tesis Doctoral, Posgrado en Ciencias Biológicas, Universidad Nacional Autónoma de México, 363 pp.
- Villalobos JL, Alvarez F (2003) Two new species of freshwater crabs of the genus *Tehuana* (Brachyura: Pseudothelphusidae) from southern Mexico. *Journal of Crustacean Biology* 23: 223–229.
- Villalobos JL, Alvarez F (2009) Los cangrejos de la familia Pseudothelphusidae (Decapoda: Brachyura: Eubrachyura) de México, con un apéndice de las especies citadas para América hasta el 2006. In: Alvarez F, Rodriguez-Almaraz G (Eds) *Crustáceos de México: Estado Actual de su Conocimiento*. Universidad Autonoma de Nuevo Leon, Secretaría de Educación Pública, Monterrey, Nuevo Leon, México, 239–300.
- Villalobos JL, Alvarez F (2010) Phylogenetic analysis of the Mexican freshwater crabs of the tribe Pseudothelphusini (Decapoda: Brachyura: Pseudothelphusidae). *Zoological Journal of the Linnean Society* 160: 457–481. doi: 10.1111/j.1096-3642.2009.00606.x
- Villalobos JL, Alvarez F (2013) Two new genera and three new species of freshwater crabs (Crustacea: Pseudothelphusidae: Potamocarcinini) from Chiapas, Mexico. *Zootaxa* 3599: 457–470. doi: 10.11646/zootaxa.3599.5.3
- Villalobos JL, Cantú A, Lira E (1993) Los crustáceos de agua dulce de México. In: Gio-Argez R, López-Ochoterena E (Eds) *Diversidad Biológica en México*. Revista de la Sociedad Mexicana de Historia Natural, Volumen Especial XLIV, 267–290.

- Villalobos JL, García AJ, Velázquez E (2010) A new species of freshwater crab of the genus *Odonthelphusa* Rodríguez, 1982 (Crustacea: Decapoda: Pseudothelphusidae) from Chiapas, Mexico. *Zootaxa* 2414: 52–58.
- Yeo CJ, Ng PKL, Cumberlidge N, Magalhães C, Daniels SR, Campos MR (2008) Global diversity of crabs (Crustacea: Decapoda: Brachyura) in freshwater. *Hydrobiologia* 595:275–286. doi: 10.1007/s10750-007-9023-3